

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number  
**WO 02/38727 A2**

(51) International Patent Classification<sup>7</sup>: **C12N**

HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(21) International Application Number: **PCT/NL01/00804**

(22) International Filing Date:  
5 November 2001 (05.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
00203911.3 9 November 2000 (09.11.2000) EP

(71) Applicant (for all designated States except US): **PLANT RESEARCH INTERNATIONAL B.V. [NL/NL]**; Droeendaalsesteeg 1, NL-6708 PB Wageningen (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VAN ENCKE-VORT, Leonora, Johanna, Gertruda [NL/NL]**; Cort van der Lindenstraat 35, NL-6702 BD Wageningen (NL). **JACOBSEN, Evert [NL/NL]**; Rizema Bosweg 77, NL-6708 DK Wageningen (NL). **STIEKEMA, Willem, Johannes [NL/NL]**; Leonard Roggeveenstraat 21, NL-6708 NB Wageningen (NL). **PEREIRA, Andy [NL/NL]**; Indira Gandhi Singel 14, NL-6717 GG EDE (NL).

(74) Agent: **DE HOOP, Eric**; Octrooibureau Vriesendorp & Gaade, P.O. Box 266, NL-2501 AW The Hague (NL).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **DNA SEQUENCES ENCODING PROTEINS CONFERRING PHYTOPHTHORA INFESTANS RESISTANCE ON PLANTS**

(57) Abstract: Genomic sequences encoding *Phytophthora infestans* resistance proteins are provided herein. Specifically, sequences from potato required for *P. infestans* resistance have been cloned and sequence provided, together with the encoded amino acid sequence. DNA encoding the amino acid sequence or amino acid sequences showing a significant degree of homology thereto may be introduced into plant cells and the encoded polypeptide expressed, conferring *P. infestans* resistance on plants comprising such cells and descendants thereof.

WO 02/38727 A2

BEST AVAILABLE COPY

**DNA sequences encoding proteins conferring *Phytophthora infestans* resistance on plants****FIELD OF THE INVENTION**

5        This invention relates to methods and materials for improved plant disease resistance. In particular the present invention relates to nucleic acid sequences required for resistance of potato to *Phytophthora infestans*, recombinant polynucleotide molecules containing the sequences, and uses thereof to transform plants, especially plants of the family *Solanaceae* to make them more resistant to *Phytophthora* species.

10

**BACKGROUND OF THE INVENTION**

      The oomycete pathogen *Phytophthora infestans*, is worldwide the main disease of the potato crop causing late blight that results in major losses of crop yield and quality. *P. infestans* infects  
15        plants of commercial importance like potato and tomato, that therefore require regular chemical control. Monogenic *R* genes have been introduced from the hexaploid Mexican wild species *Solanum demissum* into the cultivated tetraploid potato cultivars (Wastie, 1991). These race specific *R* genes did not provide durable field resistance because of the rapid evolution of new virulent races of the fungus that circumvent these *R* gene mediated resistances. Characteristic  
20        for *R* gene mediated resistance reactions is the hypersensitive response (HR) leading to local cell death causing necrotic spots at the site of attempted infection. Genetic analysis showed that activation of HR is highly specific and induced upon recognition by a specific *R* gene product and a corresponding avirulence gene product in the pathogen (Hammond-Kosack and Jones, 1997).

25        The *R* gene mediated resistance from wild *Solanum* species can show partial resistance or an intermediate HR response when crossed to different *S. tuberosum* backgrounds (Graham, 1963; Toxopeus, 1958). The HR lesions can vary in size depending on the backcross parent used, indicating that other genes influence the *R* gene resistance reaction. Minor *S. tuberosum* or *S. demissum* genes have been characterized to influence or even suppress *R* gene expression  
30        (El-Kharbotly *et al.*, 1996b). QTL mapping in *S. tuberosum* populations segregating for partial *P. infestans* resistance, identified 19 QTLs on 13 chromosomal regions (Leonards-Schippers *et al.*, 1994), with one QTL on chromosome 5 near the *P. infestans* resistance locus *R1* also linked to QTLs for maturity and vigor (Collins *et al.*, 1999). These QTLs on chromosome 5 very likely represent minor genes that play a role in both *R* gene mediated HR resistance responses

and developmental processes which indirectly influence the resistance response. Additionally, this chromosome region also contains several other resistance loci with specificity to different pathogens like the PVX virus (Ritter *et al.*, 1991) and potato cyst nematodes (Kreike *et al.*, 1994; Rouppe van der Voort *et al.*, 1998).

5       The cloning of *R* genes that mediate gene-for-gene type resistance to bacterial, fungal, oomycete, viral, and nematode pathogens has so far identified 5 classes of genes based on common characteristics including nucleotide binding sites, leucine-rich repeats, transmembrane domains and serine/threonine protein kinases (Hammond-Kosack and Jones, 1997). Genetic mapping and sequence analysis showed frequent clustering of *R* genes with different resistance  
10       specificities at complex loci (Jia *et al.*, 1997; Parniske *et al.*, 1997). Despite these insights into *R* gene structure their function can not be predicted from sequence alone and functional tests are required to determine their role in resistance (Parker *et al.*, 1996).

A few *R* gene signal transduction components have been identified by mutation (reviewed in Innes, 1998). These analyses have helped identify genes that are required for the barley  
15       powdery mildew mediated *Mla-12* resistance (*rar-1* and *rar-2*; Jørgensen, 1996), the tomato *Pseudomonas syringae* pv *tomato* resistance gene *Pto* (*Prf*, Salmeron *et al.*, 1996) and *Pti*; (Zhou *et al.*, 1995) and for the tomato *Cf-9* (*rcr-1* and *rcr-2*; Hammond-Kosack and Jones, 1994) and *Cf-2* mediated *Cladosporium fulvum* resistance reactions (*rcr-3*; Jones *et al.*, 1999). Extensive mutant screens in *Arabidopsis* identified a number of genes involved in plant  
20       pathogen interactions, *ndr1* (Century *et al.*, 1995), *eds1* (Parker *et al.*, 1996), *pad1*, *pad2*, *pad3* and *pad4* (Glazebrook *et al.*, 1996) and *pbs1*, *pbs2* and *pbs3* (Warren *et al.*, 1999). Most of these mutations affect the function of a subset of *R* genes (Aarts *et al.*, 1998) or only combinations of double mutations significantly decrease *R* gene resistance (Glazebrook *et al.*, 1997; Warren *et al.*, 1999; McDowell *et al.*, 2000). This indicates the occurrence of different  
25       signaling pathways for resistance reactions that are also partially redundant.

Transposon tagging is an established tool in plants for the identification of genes that display a mutant phenotype when their function is disrupted. Transposons have been introduced from maize and successfully used for tagging in many heterologous plants like *Arabidopsis* (Aarts  
*et al.*, 1993), *petunia* (Chuck *et al.*, 1993), *tobacco* (Whitham *et al.*, 1994), *tomato* (Jones *et al.*,  
30       1994) and *flax* (Lawrence *et al.*, 1995). In these self-fertilizing plant species random tagging strategies (*Arabidopsis*, *petunia*) by screening large selfed populations for mutants or targeted tagging of specific genes (*tobacco*, *tomato*, *flax*) were applied. By self- or test- crossing, large populations were produced for the direct screening of possible transposon tagged mutants. By using selectable markers like kanamycin (Baker *et al.*, 1987) or hygromycin (Rommens *et al.*,

1992), selection of excision events at the cellular level has been feasible and in combination with effective in vitro selection and somatic propagation procedures can facilitate the production of large numbers of transposon insertion mutants.

## 5 SUMMARY OF THE INVENTION

The present invention provides an isolated DNA sequence which encodes a protein having the amino acid sequence given in SEQ ID NO:2 or a functionally homologous protein having an amino acid sequence showing an identity of at least 80% to SEQ ID NO:2, which protein  
10 confers *Phytophthora infestans* resistance on plants.

Preferably, the DNA sequence encodes a protein having an amino acid sequence showing an identity of at least 85%, or even 90%, to SEQ ID NO:2.

More preferably, the DNA sequence encodes the amino acid sequence given in SEQ ID NO:2, in which case the DNA sequence comprises the nucleotide sequence given in SEQ ID  
15 NO:1.

More general, the invention provides the DNA sequence selected from the group consisting of:

- a) the DNA sequence given in SEQ ID NO:1 and its complementary strand, and
- b) DNA sequences hybridizing to the sequences in (a) under stringent hybridization  
20 conditions.

In particular the DNA sequence comprises the nucleotide sequence of nucleotides 20 to 1053 of SEQ ID NO:1, which is the coding sequence for the protein having the amino acid sequence given in SEQ ID NO:2. Said nucleotide sequence contains an intron, nucleotides 584 to 726. Accordingly, the actual DNA sequence encoding the protein of SEQ ID NO:2  
25 comprises nucleotides 20 to 583 and 727 to 1053 of SEQ ID NO:1.

In a further aspect the invention provides a protein having the amino acid sequence given in SEQ ID NO:2 or a functionally homologous protein having an amino acid sequence showing an identity of at least 80% to SEQ ID NO:2, which protein confers *Phytophthora infestans* resistance on plants.

30 In another aspect the invention provides a recombinant vector comprising a DNA sequence as defined above under control of an appropriate promoter and regulatory elements for expression in a host cell.

In still another aspect the invention discloses the use of the present DNA sequence or recombinant vector for the production of a transgenic plant.

Further the invention provides a host cell, preferably a plant cell, comprising the present DNA sequence or recombinant vector.

Also provided is a plant or any part thereof comprising such plant cell, and seed, selfed or hybrid progeny or descendant of such a plant, or any part thereof.

5 The invention also provides a method of conferring *Phytophthora infestans* resistance on a plant, comprising the steps of

- i) introducing a DNA sequence as defined above or a recombinant vector as defined above into a cell of the plant or an ancestor thereof,
- ii) regenerating plants from the obtained transgenic cells, and
- 10 iii) selecting plants exhibiting *P. infestans* resistance.

A further aspect of the invention is the provision of oligonucleotide probes that comprise a sequence of nucleotides of SEQ ID 1, or a mutant, derivative or allele thereof, capable of detecting the pathogen resistance gene or functional equivalents thereof in plants of the  
15 family *Solanaceae* and the use of the probes to isolate DNA sequences encoding a pathogen resistance gene or a functional equivalent thereof.

Using the sequence SEQ ID 1 facilitates the isolation of homologous genes from related and unrelated hosts to obtain genes, which protect host plants against related and unrelated pathogens.

20 A further aspect of the invention is the identification of proteins that interact with constructs comprising sufficient homology to SEQ ID 2, the genes thereof can be used to provide plant cells that are resistant to pathogens. One way is by identification of interacting proteins by the yeast two-hybrid system that are then involved in the signal transduction of the resistance response.

25 A further aspect of the invention is the construction of hybrid proteins comprising SEQ ID 2 or DNA isolates of sufficient homology, with other proteins that can be used as effector molecules. One way is by making hybrids with different leucine rich repeat fragments from various plants or synthetically produced *in vitro*, that can interact with different pathogen or inducer effector molecules. These effectors can also be chemically  
30 produced and by application to a plant containing the hybrid construct can induce the signal transduction pathway for resistance.

**BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1: Schematic drawing of pHPT::Ds-Kan showing positions of primer 1 (p1, GCG CGT TCA AAA GTC GCC TA), primer 2 (p2, GTC AAG CAC TTC CGG AAT CG) and  
 5 *Pst*I restriction sites. Abbreviations: LB = left border, RB = right border, pNOS = nopaline synthase promoter, NPT II = neomycin phosphotransferase gene, HPT II = hygromycin phosphotransferase gene.

Figure 2: *Pst*I restriction of genomic DNA hybridized to NOS promoter probe to select for  
 10 presence of full donor site (FDS = 4.0 kb), empty donor site (EDS = 2.3 kb), *Ac* T-DNA construct (3.5 kb) and *Ds* re-insertion sites in the R1Ds/r-; *Ac*- selected seedlings EE96-4311-37 (lane 1), EE96-4312-43 (lane 5) EE96-4312-49 (lane 9) and HygR protoplast regenerants from EE96-4311-37 (lane 2,3 and 4), EE96-4312-43 (lane 6, 7 and 8), EE96-4312-49 (lane 10, 11, 12 and 13), EE96-4311-15 (lane 13, 15), EE96-4312-05 (lane 14),  
 15 EE96-4312-76 (lane 16) and EE96-4312-06 (lane 17).

Figure 3: Reaction phenotypes observed on different genotypes after inoculation of detached leaves with *P. infestans* race 0. a) TM17-2, susceptible parent; b) detail of sporulation on TM17-2; c) HRPR 836; d) HRPR 1587 showing both the *R1* type HR response and necrotic regions with sporulation; e) detail of HR spot on HRPR 1587; f) detail of sporulation on the necrotic region of HRPR 1587; g) necrotic regions on R1Ds/r-; *Ac*- seedling EE96-4312-03, minor sporulation was detected in such regions; h) clear colonization on variant 1000; i) detail of sporulation on variant 1000.

25 Figure 4: *Hind*III digested genomic DNA hybridized to the 5' *Ac* probe (A) or the internal *Ac* probe (B). Lane 1 shows the 1.6-kb marker hybridization. The *R1* resistant crossing parent J91-6167-2 (lane 2a and b) and the susceptible crossing parent 87-1024-2 (lane 3a and b) contain both no *Ac* or *Ds* elements. The primary transformant Ds416 contained two *Ds* T-DNA loci (lane 4a), Ds53-34 inherited both *Ds* T-DNA loci (lane 5a) as did EE96-4312-28 (lane 7a). EE96-4312-28 inherited from TM17-2 (lane 6a) the *Ac* element. In mutant 487 (lane 8a) and mutant 1000 (lane 9b), both regenerated from EE96-4312-28, the *Ds* elements transposed to new positions and *Ac* seems to be missing. In TM17-2 (lane 6b) a complete *Ac* (1.6 kb internal *Hind*III fragment) and a *dAc* (2.9 kb) are present. Mutant 487

(lane 8b) inherited *dAc* as a different restriction fragment due to the insertion of *Ac* in *dAc*. In mutant 1000 (lane 9b) *Ac* got lost and only *dAc* is present.

- Figure 5: a) Schematic representation of the isolated *Ds* flanking sequences from mutant 1000 (*rpr1* and *rpr2*) and their alignment to XA21 aa 708- 1011. The large triangles represent the *Ds* positions, the intron region is a dashed line and the small black triangles represent primer positions of EE1 (5'- ACA TTG GGC ACT CTT GGA TAC A), EE2 (5'- TCT TGA TTC TGG CAT TTT CTT TG), EE3 (5'- CCT GAC ACA AAC CGA GAC ATT), EE6 (5'- AAC AAT GCC TTT CTT CTC), EE8 (5'- GCA CAT TAT CAA GTG GAA CTA CG) and EE10 (5'- CTG AGC CGT ACT CTT AAA AGA ACG). b) Amino acid alignment of Pto, Pti1, StPK-B, StPK-A and Xa21. The eleven conserved domains of a protein kinases are numbered and the conserved amino acids are marked (\*). Bold domains are specific for serine/threonine recognition. The N-glycosylation site is underlined.
- Figure 6 (A and B): Sequence of the StPK-B DNA (SEQ ID NO: 1) and StPK-B protein (SEQ ID NO: 2). StPK-B DNA sequence is the contiguous sequence of a DNA fragment obtained by isolating the *Ds* tagged gene. The sequence is part of the StPK-B gene encoding the protein kinase domain and lacks the N-terminal portion of the gene including the translation start. SEQ ID 2 is the amino acid translation of SEQ ID 1 after removing the predicted intron conserved with other gene family members.

- Figure 7: DNA sequences of StPK homologs obtained by using primers EE1 and EE2. SEQ ID 3 is the sequence of DNA fragments of StPK-A that is tagged by the transposon *Ds*. SEQ ID 4-12 are DNA fragments obtained by PCR between primers EE1 and EE2; Seq ID 4-12 represent fragments of homologous sequences StPK-C to StPK-K, in the genome.

- Figure 8: Diagram of StPK overexpression constructs used for plant transformation. The StPK-B gene with a synthetic translation initiation start, under control of the CaMV 35S promoter and Nos-terminator is cloned in a binary vector for transformation of plants using *Agrobacterium tumefaciens* strains in construct I. Construct II contains the N-terminal part of the StPK genes or other N-terminal fusions to protein domains that act as recognition domains with other effector molecules.

## DETAILED DESCRIPTION OF THE INVENTION AND EXPERIMENTS

### Development of transposon mutagenized potato plants

Diploid potato plants heterozygous for the *P. infestans* *R1* resistance gene were transformed with an *Agrobacterium* strain containing a *Ds*-transposon T-DNA construct shown in Figure 1 (Pereira *et al.*, 1992; El-Kharbotly *et al.*, 1995). Transformant Ds416 contained a *Ds* T-DNA insertion on chromosome 5 (El-Kharbotly *et al.*, 1996a), linked in repulsion phase to the previously mapped *P. infestans* *R1* resistance gene (Leonards-Schippers *et al.*, 1992). This Ds416 clone was crossed to the susceptible diploid genotypes J89-5040-2 producing offspring that enabled the selection of recombinant plants (Ds53-22 and -34) having the *R1* gene and the *Ds* T-DNA in coupling phase (18 cM) (El-Kharbotly *et al.*, 1996a). To activate the *Ds* transposon these plants were crossed with TM17-2, a diploid potato clone susceptible to *P. infestans* and transformed with the *Ac* transposon-containing T-DNA construct pMK1GBSS*Ac* (Pereira *et al.*, 1991). TM17-2 contained one functional *Ac* displaying active transposition. From the progeny of these crosses, population EE96-4311 (Ds53-22 X TM17-2; 18 seedlings) and EE96-4312 (Ds53-34 X TM17-2; 96 seedlings), 47 (8 and 39) kanamycin resistant *R1* seedlings (KanR *R1*) were selected.

Plant genomic DNA was isolated from greenhouse grown leaves (Pereira and Aarts, 1998) and used for molecular analysis. Empty donor sites (EDS-PCR) indicating excision were detected in 22 of the 47 KanR *R1* seedlings as a 450-bp PCR product using specific primers (Figure 1) and confirmed by Southern blot analysis. After selection of the 22 *R1* resistant seedlings showing active *Ds* excision (*R1Ds/r-; Ac/-*), the expression of hygromycin resistance (HygR) was tested by rooting on MS30 supplemented with 10-100-mg/l hygromycin. One genotype EE96-4311-12 showed resistance by rooting on 40 mg/l hygromycin and displayed a clear EDS fragment, suggesting that screening for rooting of shoots on 40 mg/l could be used as a stringent criteria for *Ds* excision.

As most genotypes contained excision events that occurred late in shoot development these HygR cells could be selected by protoplast isolation and screening for hygromycin resistance to select independent excision events. Protoplasts were isolated from 4-week-old *in vitro* grown shoots (Uijtewaal *et al.*, 1987), re-suspended in culture medium TM2G (Wolters *et al.*, 1991) to a final concentration of 500,000 pp/ml and diluted weekly with fresh medium. The regenerating calli were progressively transferred to callus growth medium, shoot induction medium and finally maintained on shoot elongation medium until regenerated plants could be harvested (Mattheij *et al.*, 1992). In separate experiments to



select specifically for protoplast regenerants with excision events, 10 mg/l hygromycin was added to the callus growth medium 14 days after protoplast isolation, then increased to 20 mg/l on day 21 and maintained at this level.

- Table 1 gives an overview of protoplast regeneration data. From parental clone Ds53-34, control EE96-4312-21 and selected R1Ds/r-; Ac/- seedlings about 50 regenerating shoots were tested for their rooting ability on MS30 with 40 mg/l hygromycin. As expected, the parent Ds53-34 and control EE96-4312-21 produced no HygR protoplast regenerants whereas EE96-4311-12 gave 45% HygR protoplast regenerants confirming early excision. The other 14 good performing R1Ds/r-; Ac/- plants showed regeneration of 4 to 33%<sup>a</sup> of HygR shoots indicating excision of *Ds* from its original T-DNA location. The use of hygromycin selection during callus culture and regeneration of shoots increased the recovery of HygR regenerants 3.8 times. A total of 1973 HygR regenerants were obtained from different selection experiments and transferred to the greenhouse.
- Table 1: Selection of excision events after protoplast regeneration with and without hygromycin selection. Number of calli, shoots and selected hygromycin resistant (HygR) regenerants for parents Ds53-22 and Ds53-34; control EE96-4312-21 (R1Ds/r-; -/-) and 22-selected R1Ds/r-; Ac/- genotypes from the seedling populations EE96-4311 and EE96-4312.
- <sup>a</sup> low due to infection.

5	Genotype	No selection			Hygromycin selection		
		During protoplast regeneration			During protoplast regeneration		
		Calli	Shoots	HygR	Calli	Shoots	HygR
10	Ds53-22	10	0		47	0	
	Ds53-34	100	45	0	134	1	0
	EE96-4312-21	100	21	0	900	10	0
	EE96-4311-08	0 <sup>a</sup>			0 <sup>a</sup>		
15	EE96-4311-12	100	49	22	1000	198	98
	EE96-4311-15	300	82	11	800	160	101
	EE96-4312-03	100	23	3	1000	166	83
	EE96-4312-05	100	29	8	1000	198	121
20	EE96-4312-06	100	6 <sup>a</sup>	2	1000	205	139
	EE96-4312-14	100	70	15	1000	208	118
	EE96-4312-23	100	51	2	1000	211	91
	EE96-4312-27	10	0		10	0	
	EE96-4312-28	100	47	7	1000	143	82
25	EE96-4312-30	100	0		419	0	
	EE96-4312-31	100	30	2	570	21	4
	EE96-4312-37	100	52	2	1000	248	92
	EE96-4312-40	33	2	0	67	0	
	EE96-4312-43	100	45	8	650	207	101
30	EE96-4312-46	14	9	0	103	0	
	EE96-4312-49	100	48	3	1000	206	109
	EE96-4312-52	3	3	0	1	1	1
	EE96-4312-60	100	52	7	1000	203	93
	EE96-4312-63	100	50	7	1000	203	130
35	EE96-4312-76	0	0		24	0	
	EE96-4312-89	100	49	4	274	41	19
Total			691	103		2619	1382

40

To analyze *Ds* excision in the HygR protoplast regenerants Southern blot hybridization was performed on a subset of selected R1Ds/r-; Ac/- seedlings and some of their HygR protoplast regenerants (Figure 2). Plant DNA was restricted with *Pst*I and the blots hybridized to probes derived from the NOS promoter fragment that revealed the *Ac* T-DNA and the *Ds* transposon. The R1Ds/r-; Ac/- seedlings used for protoplast isolation all displayed two *Pst*I fragments, respectively 4.0- and 3.5-kb, corresponding to respectively the *Ds* T-DNA and the *Ac* T-DNA constructs. Faintly visible fragments of 2.3-kb were also detected that correspond to a low amount of EDS fragments present in these seedlings. All

45

HygR protoplast regenerants showed a strong hybridizing EDS fragment indicating early or repeated excision of *Ds* corresponding to the high level of hygromycin resistance for which these plants were selected. The original *Ds* parent had two copies of *Ds* at one locus. Full donor site fragments were detected in most of the HygR protoplast regenerants which indicates that one of the two *Ds*'s was not excised. Three plants shown in Fig 2, showed a complete EDS indicating that excision occurred in the initial protoplast. Most HygR regenerants showed clear *Ds* re-insertion fragments varying from 1 to 8 new positions per individual HygR regenerant. Regenerants from a single seedling showed different re-insertion patterns, indicating that they originated from independent transposition events and confirmed that most selected HygR regenerants originate from independent transposition events.

The somatic selection of *Ds* transpositions from individual cells facilitated the production of a large population of shoots with independent *Ds* excision events. The HygR protoplast regenerants potentially represent about 2000 independent *Ds* insertions. This number of *Ds* insertion mutations should be enough for the isolation of tagged mutants involved in *R1* resistance. The somatic selection of *Ds* transposition and the rapid production of independent plants containing these transpositions, facilitates the production of large tagging populations needed for the transposon mutagenesis of selected genes. This is particularly suitable for the mutagenesis of genes in heterozygous crops like potato.

20

#### Screen for *R1* type HR resistance variants in the *Ds* tagged population

The transposon mutagenized population was suitable for the isolation of mutations in defense related genes causing an altered reaction to *P. infestans*. By using a suitable screen quantitative changes towards susceptibility were possible to be identified. Race specific resistance *Cf* genes in tomato have shown a semidominant phenotype if screened in a quantitative manner (Hammond-Kosack and Jones, 1994). Chromosome 5 in potato is known to contain many resistance components (Leonards-Schippers *et al.*, 1994) that are probably in a heterozygous state as seen from segregation of minor factors. These loci could probably be efficiently mutagenized due to active linked transposition of *Ds* near *R1*.

To prepare the inoculum for screening (El-Kharbotly *et al.*, 1994), *P. infestans* race 0 (89148-09) was grown on rye agar medium (with 20 mg/l sucrose). The sporangiospores were washed with 10-15 ml cold tap water (4°C) and the resulting suspension used to inoculate 10 Bintje tuber slices (1 cm thickness). The newly formed sporangiospores were washed and again used to inoculate 20-50 tuber slices of Bintje in order to obtain 1-2 l of

sporangiospore solution. This solution was diluted to contain at least 2000 spores/ml to use for plant inoculation.

The 1973 hygromycin resistant protoplast regenerant (HRPR's) plants to be tested were periodically brought in batches to the greenhouse. After 6-10 weeks growth two leaves of  
5 each HRPR plant were harvested, placed in columns of water absorbent substrate, and put in containers (46 x 31 x 8 cm) closed with transparent covers. In every container two leaves of 10 HRPR plants and a leaf of the susceptible control (Bintje or TM17-2) were tested. In each experiment 15-30 containers were used so that 150-300 plants could be tested in parallel, with Ds53-22, Ds53-34 and TM17-2 always tested as additional controls. Each leaf  
10 in the experiments was sprayed with about 5-10 ml of the sporangiospore solution containing 10,000-50,000 sporangiospores. After 5 days in high humidity at 16°C, all leaves were evaluated for the development of *P. infestans* infection symptoms and at day 6 a second evaluation for disease symptoms was performed. When development of symptoms occurred the leaves were kept for an additional 2 days for a microscopic examination of the  
15 disease development.

The susceptible parent and control cultivar Bintje always showed distinctive colonization and abundant sporulation on day 5-6 (Figure 3a and 3b). In contrast the resistant parents Ds53-22, Ds53-34 and most of the analyzed HRPR's always displayed characteristic *R1* type HR spots upon infection. The phenotype of HRPR 836 was distinctly susceptible with  
20 colonization and sporulation over large leaf areas (Figure 3c). Other HRPR's sometimes showed larger necrotic regions indicating colonization of the leaves (Figure 3d). When this colonization resulted in sporulation (Figure 3f) the HRPR was scored as a potentially susceptible *R1* variant, although necrotic spots were additionally visible on the green parts (Figure 3e) indicating at least a partial HR activation. In this first round of screening 33  
25 putative susceptible variants, derived from 10 R1Ds/r-; Ac/- seedlings were selected (Table 2). Re-inoculation tests of newly grown leaves of the selected variants confirmed the susceptible reactions for 9 variants.

Table 2: Primary screen for mutants with an altered *RI* type HR resistance response.

	RIDs/r-/Ac/- Seedling	total # selected HRPR's	# HRPR's tested with <i>P. infestans</i> race 0	# HRPR's Variant	Variant plant #	Ploidy level
5	EE96-4311-12	126	72	1	702	4x
	EE96-4311-15	112	81	2	35	2x
					994	2x
	EE96-4312-03	86	63	2	1515*	2x
					1921	nd
	EE96-4312-05	129	84	2	836*	4x
					842	4x
10	EE96-4312-06	243	188			
	EE96-4312-14	195	168	2	925	4x
					1587	2x
	EE96-4312-23	93	82			
	EE96-4312-27	5	0			
	EE96-4312-28	89	71	7	487*	2x
					998	2x
					999	4x
					1000	2x
					1001	4x
					1005	4x
					1357	2x
15	EE96-4312-31	7	4			
	EE96-4312-37	134	120	4	151*	4x
					510*	4x
					524	2x
					551	4x
	EE96-4312-43	109	91	6	570*	4x
					688*	4x
					1528	4x
					561	x-4x
					562	x-4x
					574	4x
	EE96-4312-49	152	111	6	600	4x
					601	4x
					633*	2x
					1050	2x
					1055*	4x
					1073	4x
	EE96-4312-52	1	1			
20	EE96-4312-60	134	112	1	667	4x
	EE96-4312-63	168	155			
	EE96-4312-76	83	65			
	EE96-4312-89	107	96			
	Total	1973	1564	33		
25	# = number, nd = not determined, * variants with confirmed susceptible reaction after re-inoculation.					

Selected genotypes were transferred from the greenhouse to *in vitro* for propagation to obtain 10 or 35 cuttings of each variant and these were transferred again to the greenhouse for a replicated re-testing of the *P. infestans* R1 resistance. From the first set of 33 R1 variants, ploidy level analysis enabled the identification of plants with chromosomal anomalies that were potentially somaclonal variants. All the diploid variants together with those with a reproducible susceptible phenotype and the corresponding 9 parental seedlings were used for a secondary quantitative phenotypic analysis. After *P. infestans* inoculation on two leaves of each plant, the developing symptoms were carefully evaluated and followed microscopically when necessary to detect sporulation (Table 3).

Table 3: Qualitative and quantitative analysis of the resistance reaction and disease development on selected variants compared to parental controls.

	# plants	% leaves with HR	% leaves with <20% necrosis and sporulation	% leaves with 20-100% necrosis, colonization sporulation (max % leaf area covered)	& Dead (Rotten)
Parents					
Ds53-34	10	100	0	0	0
TM17-2	10	0	0	100 (100)	0
R1Ds/r-; Ac/- Seedlings					
EE96-4312-76	10	100	0	0	0
EE96-4312-43	9	94	6	0	0
EE96-4312-37	19	87	13	0	0
EE96-4312-28	20	75	23	2 (35)	0
EE96-4312-49	10	75	20	5 (25)	0
EE96-4312-05	20	75	15	5 (40)	5
EE96-4312-03	21	71	19	10 (60)	0
EE96-4311-15	20	60	35	5 (50)	0
EE96-4312-14	19	47	18	32 (100)	3
Variants					
EE96-4312-43 570	9	100	0	0	0
688	7	100	0	0	0
EE96-4312-37 510	10	95	5	0	0
524	10	90	5	0	5
EE96-4312-28 487	35	41	43	16 (100)	0
998	34	60	38	2 (35)	0
1000	35	16	34	50 (100)	0
1357	34	63	24	9 (70)	4
EE96-4312-49 601	10	60	30	10 (60)	0
633	10	70	25	5 (60)	0
1050	9	78	17	5 (45)	0
1055	10	80	10	5 (50)	5
EE96-4312-05 836	34	7	12	68 (100)	13
EE96-4312-03 1515	31	76	16	8 (50)	0
EE96-4311-15 35	14	82	0	7 (50)	11
994	8	44	0	56 (100)	0
EE96-4312-14 1587	1	50	0	50 (60)	0

# = number, HR = hypersensitive response

The *RL*-resistant parental plant Ds53-34 always showed the complete *RI* type HR response, with small necrotic spots on inoculated leaves. The *RI* resistant progeny of Ds53-22 and Ds53-34 (EE96-4311-15 and EE96-4312-03, 05, 14, 28, 37, 43, 49 and 76) displayed an intermediate resistance phenotype (Table 3). With the exception of seedling 5 EE96-4312-76, all other seedlings showed on several leaves (6-35%) necrotic spots that developed into necrotic regions covering 5 to 20% of the leaf area (Figure 3g). Microscopic examination revealed very little sporulation in these regions indicating minor escape of *P. infestans* from the normal *RI* type HR response. In a few leaves the necrotic region covered almost 100% of the leaf area and colonization with sporulation was observed indicating 10 susceptibility of the leaf and escape from the *RI* type HR resistance response. Seedling EE96-4312-14 showed in this analysis only in 47% of the leaves a clear *RI* type HR response. However, from the 168 HRPR's derived from this seedling and tested in the first screening for *RI* resistances only 2 were selected as putative variants (Table 2). This indicates that the intermediate phenotype for this and other parental seedlings did not result 15 in an overestimation of putative variants in the first screening.

The re-evaluation of the resistance response reaction for the variants 487, 1000, 836 and 994 showed a clear deviation in phenotype when compared to the parental seedlings. Variant 1000 showed colonization and sporulation on 50% of the leaves and this clearly resembled the TM17-2 *P. infestans* susceptible parental phenotype (Figure 3h and 3i). Only 16% of the 20 variant 1000 leaves showed the normal *RI* type HR resistance response. In variant 487 the *RI* type HR resistance response was clearly detected in only 41% of the inoculated leaves. On 16% of the leaves necrotic regions covered over 20% of the leaf area and colonization and sporulation was detected indicating a weak susceptible *RI* variant.

The susceptible phenotype of variant 836 in the first screening of the HRPR population 25 (Figure 3c) was repeatable in this analyses but quick senescence of the leaves, resulting in softening and rotting, suggested other causes for the observed susceptible phenotype. Variant 994 showed a striking phenotype as in every plant the youngest leaf showed colonization with sporulation, combined with leaf softening and rotting. The second oldest leaf analyzed always showed a normal *RI* type HR resistance phenotype. The variants 836 and 994 30 therefore displayed a more susceptible reaction due to interaction with early senescence and were therefore not considered as variants in the expression of *RI* type resistance. Re-evaluation of the resistance phenotype of the variants 570, 688, 510, 524, 633, 1050, 1055, 1515 and 35 did not reveal any quantitative difference when compared to the parental seedlings and were not regarded as mutants in the *RI* resistance reaction.

### Molecular analysis of the tagged mutants

To examine the causal relationship between the *Ds* insertion sites and the observed phenotype mutant 1000 was characterized by Southern blot hybridization. Genomic DNA from appropriate genotypes was restricted with *Hind*III and the blots hybridized to a 5' *Ac* probe to determine the presence and positions of the *Ac* and *Ds* elements in the different genotypes. Since *Ds* is derived from *Ac*, the 5' *Ac* probe also identified the *Ds* element (Pereira *et al.*, 1992). Additional hybridization of the same blots with an internal *Ac* probe revealed the presence and position of *Ac*. In the different genotypes the *Ds* and *Ac* insertions were identified by specific *Hind*III fragments (Figure 4a and 4b). New positions of the *Ds* elements confirmed that both mutants (from same seedling parent) were derived from different *Ds* transposition events in EE96-4312-28 during protoplast regeneration (Fig 4a). These hybridizations also revealed that mutant 1000 had lost the *Ac* element and was therefore a stable mutant.

To analyze the sites of *Ds* insertion, flanking DNA of *Ds* insertions was isolated by inverse PCR (IPCR ;Triglia *et al.*, 1998). Plant genomic DNA, was restricted with *Hae*III, self-ligated and restricted with *Bam*HI and *Bg*II. *Bg*II restriction prevents the amplification of the *Ds* transposon flanking sequences in the original T-DNA construct. To obtain additional and longer 5' flanking sequences a second restriction combination was used, in which genomic DNA was restricted with *Msc*I followed by *Hind*III and after ligation linearized by *Cla*I. Primer 5'-CGG GAT GAT CCC GTT TCG TT (*Ac* position 197-216) and primer 5'-GAT AAC GGT CGG TAC GGG AT (*Ac* position 44 - 35) were used to amplify the 5' *Ds*/*Ac* flanking sequences. After a hot start (10 min 94°C), 35 PCR cycles (1 min 94°C, 1 min 60°C, 2 min 72°C) resulted in the amplification of *Ac* and *Ds* 5' flanking sequences.

Thermal asymmetric interlaced (TAIL) PCR (Liu and Whittier, 1995) was used to obtain additional *Ds* transposon flanking sequences. Sets of nested primers designed on the 5'- and the 3' site of the *Ac* transposon (Tsugeki *et al.*, 1996; Ds5-1, 5-2, 5-3, 5-4 and Ds3-1, 3-2, 3-3, 3-4) were combined with 4 different degenerated primers (AD1 to 4; Liu and Whittier, 1995) or two other degenerate primers (Tsugeki *et al.*, 1996; renamed AD5 and 6). The three step PCR reactions were performed as described (Tsugeki *et al.*, 1996). Primers AD3, AD5 and AD6 with *Ac*/*Ds* 5' primers and primers AD5 and AD6 with *Ac*/*Ds* 3' primers produced specific PCR fragments.



The IPCR and TAIL-PCR products were separated on a 1% TBE agarose gel to determine the number and size of fragments. After phenol:chloroform extraction and isopropanol precipitation, the PCR products were cloned in pGEM T easy vector (Promega Corporation). For each sample three clones were sequenced using an automated ABI 373 DNA sequencer. The obtained *Ds* flanking sequences were compared to known sequences by BlastN and BlastX homology searches (Altschul *et al.*, 1997) in the public databases.

In mutant 1000 two different *Ds* insertions were identified by BlastX searches, each with homology to a leucine rich repeat containing protein kinase from *Oryza longistaminata* (Tarchini *et al.*, 2000) and a receptor protein kinase-like protein from *Arabidopsis thaliana* (BAC clone F13I12). These sequences were both identified due to their homology to the serine/threonine kinase domain of the *Xanthomonas* resistance gene *Xa21* isolated from *O. longistaminata* (Song *et al.*, 1995). Additional *Ds* flanking sequences were isolated using TAIL-PCR (540 bp; Fig 5a). Combining the 5' 288 bp and the 3' 540 bp flanking sequence for this *Ds* insertion revealed the expected 8-bp target site duplication. The sequence flanking this *Ds* insertion showed 50% protein identity to XA21. For the second *Ds* insertions the IPCR and TAIL-PCR only extended the 5' flanking sequence from 496 to 1309 bp (Fig 5a). This sequence covered a complete serine/threonine protein kinase ORF (nucleotides 20 to 583 and 727 to 1053) with 44% identity to the serine/threonine protein kinase domain of XA21 including the conserved intron position, nucleotides 584 to 726 (Fig 5a). All eleven protein kinase specific domains with conserved features were present (Fig 5b) as well as all the 14 conserved amino acids (Hanks *et al.*, 1988). Domain VI (consensus DLKPEN) and domain VIII (consensus G(T/S)XX(Y/F)XAPE) are indicative of serine/threonine specificity (Hanks *et al.*, 1988). The two *Ds* insertion loci displayed 84% identity at the protein level while in the intron region they showed only 52% nucleotide identity. This was a clear indication that the isolated *Ds* flanking *Solanum tuberosum* protein kinase (StPK) represented two distinct *Ds* tagged loci in mutant 1000, StPK-A and StPK-B. Fig. 6(A) shows the nucleotide sequence of StPK-B (SEQ ID NO: 1) and Fig. 6(B) shows the deduced amino acid sequence of StPK-B (SEQ ID NO: 2).

Although the protein identity was less than 50%, all characteristic protein kinase domains and conserved amino acids were present in the potato insertion loci StPK-A and StPK-B (except for domain X and XI in StPK-A), including the intron position at exactly the same position in the serine/threonine specific domain VIII. Therefore, it is probable that these serine/threonine protein kinases are similarly functional in the signal transduction pathway leading to *P. infestans* resistance and perhaps other pathogens.

Surprisingly, the StPKs are more homologous to the protein kinase domain of the rice resistance gene *Xa21* than to the earlier identified *Solanaceous* tomato resistance genes *Pto* (Martin *et al.*, 1993) and *Pti* (Zhou *et al.*, 1995). More surprisingly, since a homologue of *Pto* was mapped to the *R1* chromosomal 5 area (Leister *et al.*, 1996). These tomato

5 serine/threonine kinases are functional in the signal transduction pathway leading to a hypersensitive response reaction upon infection with *Pseudomonas syringae* pv. tomato strains expressing the avirulence gene *avrPto* (Zhou *et al.*, 1997). In *Xa21*, other rice homologs (Tarchini *et al.*, 2000) and in the StPK-A and StPK-B the conserved intron position in domain VIII indicates a conserved gene family among monocots and dicots. No

10 intron position was identified in the tomato serine/threonine kinases *Pto* and *Pti1*. Among the 11 kinase specific domains only minor differences were observed between the potato kinases and *Xa21* on one hand and *Pto* and *Pti1* on the other hand (Fig 5b). But overall amino acid homology determined that the potato sequences were more related to *Xa21* than to the tomato kinases *Pto* and *Pti1*. Domain IV, with no general consensus, showed a high

15 homology between *Xa21* and the potato sequences while *Pto* and *Pti1* contained different amino acids in this area. Whether this difference determines a clear difference in function or signaling pathway for these kinases needs to be studied.

#### Identification of potato protein kinase (StPK) homologs

20

To characterize the structure of the StPK homologs in *R1* resistant and susceptible plants several sets of primers were designed and used in PCR analysis. Primers EE1 and EE2 (Fig 5a) could amplify a product of expected size of about 470 bp and a second product of about 370 bp from the *R1* resistant parent J91-6167-2, the susceptible parent 87-1024-2 and

25 several *R1* resistant and susceptible progeny (J92-6400-A1, -A2, -A3, -A4, -A5 and -A6). Sequencing the PCR products derived from J91-6167-2, 87-1024-2, J92-6400-A1 and -A4 identified 10 different StPKs (Table 4). In Fig. 7 the DNA sequences of the StPK homologs are shown (partial sequences of the PCR products), SEQ ID NO:3-12. StPK-A was not identified in any of the plants by using this primer combination. From the susceptible

30 parental clone 87-1024-2, StPK-B was isolated. Two additional StPK homologs, StPK-C and -D were identified several times in both *R1* resistant and susceptible clones. StPK-D was the 370 bp PCR product and had a deletion of 108 bp making it very likely a non-functional StPK. From the *R1* resistant plants two additional StPK homologs, StPK-E and -I were isolated and from the susceptible plants 5 additional homologs were isolated, StPK-F, -G, -

H, -J and -K (Table 4). The sequences of StPK-F and StPK-G shown in Fig. 7 are smaller than they actually are due to sequencing problems. The isolation of these many StPK homologs indicated that these serine/threonine protein kinases represent a multigene family in *S. tuberosum*. This was confirmed by DNA hybridization, since StPK-B identified over 30 hybridizing fragments in *Hind*III or *Eco*RI digested genomic DNA of a single resistant or susceptible potato clone.

Table 4: Homologs of *Solanum tuberosum* protein kinases (StPK) isolated from *R*1 resistant and susceptible clones using primers EE1 and EE2 (Fig 5a)

StPK homologue	% identity to StPK-B	R1r parent 6167-2	rr parent 1024-2	R1r progeny J92-6400-A4	rr progeny J92-6400-A1	Total Clones
StPK-B	100		1			1
StPK-C	92	2		3	2	7
StPK-D $\Delta$	82	5	3	5		13
StPK-E	91			1		1
StPK-F	91		1			1
StPK-G	86		1			1
StPK-H	90				1	1
StPK-I	89	1				1
StPK-J	91		1			1
StPK-K	94				2	2
Total		8	7	9	5	29

#### $\Delta$ deletion locus

A second set of primers, EE3 and EE6 (Fig 5a), of which primer EE6 is located downstream of the second exon of StPK-B, was specific enough to identify StPK-B in all analyzed plants. So, this StPK homologue is present in 87-1024-2 (identified with EE1 and EE2), in J91-6167-2 and in several tested *R*1 resistant and susceptible progeny of population J92-6700, including -A16 from which the tagging population was derived. The StPK-B gene is therefore independent of the *R*1 locus.

A third set of primers, EE8 and EE10 (Fig 5a), was designed on low homologous regions between StPK-A and StPK-B and specifically identified the StPK-A locus after *Bgl*III digestion of the PCR products. Analyses of all the parental genotypes used in the different crossings identified that StPK-A is present in the susceptible parent 87-1024-2 that was used to produce the starting population from which J92-6400-A16 was selected (El-Kharbotly *et al.*, 1995).

Both *Ds*-transposon insertions in mutant 1000 are loci that occur solely or also in plants that do not carry the *R1* gene. Therefore, it is very unlikely that StPK-A or StPK-B are the *R1* gene itself. The *Ds* mutagenized StPK loci were designated respectively *rpr-1* and *rpr-2* (Required for *Phytophthora infestans* resistance). Both homologs cover a complete (or  
 5 almost complete) serine/threonine protein kinase ORF with all conserved characteristics including a conserved intron position. The *Ds* insertions in *Rpr1* and *Rpr2* probably reduce their expression explaining the incomplete *R1* type HR resistance reaction in mutant 1000. Examples of such mutants that can produce a phenotype are given by mutation in one or two genes of a multigene family (Gilliland *et al.*, 1998). The mutations may also be  
 10 semidominant due to a specific structure as described due to transposon or T-DNA insertions or inversions (Bender and Fink, 1995) (English and Jones, 1998) (Stam *et al.*, 1998).

If the StPK homologs are similar to the *Xa21* gene structure with an LRR additional to the kinase domain, then in StPK-A the *Ds* insertion in the serine/threonine kinase, 46 bp upstream of the intron, would probably form a truncated LRR protein without a functional  
 15 kinase domain. This putative truncated LRR domain could possibly compete with the functional LRR-kinase genes, reducing or delaying the signal transduction to exhibit partial *P. infestans* resistance.

StPK-B contains a *Ds* insertion downstream of a serine/threonine protein kinase. For this insertion *Ds* 5' promoter activity (Rudenko *et al.*, 1994) could result in the production of an  
 20 antisense RNA. Post transcriptional gene silencing due to the formed aberrant RNA could result in a reduction of kinase activity making the signaling pathway leading to the *R1* type HR response less effective. This might explain the semi-dominant mutation leading to a mutated *R1* resistance phenotype in regenerant 1000. A delay in HR response could allow escape of the *P. infestans* from necrotic regions resulting in sporulation and further  
 25 colonization of the infected leaves.

#### Transformation of StPK gene constructs conferring resistance

The StPK-B gene fragment was isolated and incorporated in binary vectors for  
 30 transforming plants. A suitable vector construct with appropriate regulatory sequences including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other appropriate sequences. The StPK-B fragment encodes the kinase domain but lacks the N-terminal part including the translation initiation part. Suitable constructs (I) and (II) are shown in Fig 8. In one construct (I) a complete kinase encoding

gene was made with the StPK-B gene consisting of the DNA fragment shown as Seq ID 1, with a translation start coding for the methionine codon in frame with the open reading frame shown in Seq ID 2. In this construct-I the engineered StPK-B gene was cloned in between the constitutive CaMV 35S promoter and a nopaline synthase (Nos) terminator in the vector pBINPLUS (van Engelen et al., 1995). In another construct type (II) a translational fusion made between the STPK-B fragment of SEQ ID1 and a N-terminal part of the complete gene from STPK-B or homologues genes is made. These gene fusions are cloned in between the appropriate regulatory promoter and terminator sequences in pBINPLUS

The above mentioned recombinant binary vectors are possible to construct by persons skilled in the art, including the transfer into appropriate *Agrobacterium* strains and checking for their stable presence in the *Agrobacterium*. The recombinant *Agrobacterium* construct containing the StPK-B overexpression cassette are transformed into potato plants by established procedures (El-Kharbotly et al., 1995). A set of transformants are regenerated and multiplied in vitro by cutting. About 5 regenerated plantlets from each individual transformant are transferred to the greenhouse.

At about 6 weeks after transferring to the greenhouse the replicated sample of the transformants are tested by the detached leaf test for *Phytophthora infestans* resistance as described in detail above. From each plant two leaflets are taken, amounting to 10 leaf samples per individual transformant. The resistance score over the replicate samples provides a quantitative estimate of the resistance reaction and allows the selection of plants significantly resistant to infection by *Phytophthora*.

## References

- Aarts, M.G.M., Dirkse, W.G., Stiekema, W.J. and Pereira, A. (1993) Transposon tagging of a male sterility gene in *Arabidopsis*. *Nature* 363, 715-717.
- Aarts, N., Metz, M., Holub, E., Staskawicz, B.J., Daniels, M.J. and Parker, J.E. (1998) Different requirements for *EDSI* and *NDR1* by disease resistance genes define at least two *R* gene-mediated signaling pathways in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America* 95, 10306-10311.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25, 3389-3402.

- Baker, B., Coupland, G., Fedoroff, N., Starlinger, P. and Schell, J. (1987) Phenotypic assay for excision of the maize controlling element, *Ac* in tobacco. *EMBO Journal* 6, 1547-1554.
- 5 Bender, J. and Fink, G.R. (1995) Epigenetic control of an endogenous gene family is revealed by a novel blue fluorescent mutant of *Arabidopsis*. *Cell* 83, 725-734.
- Century, K.S., Holub, E.B. and Staskawicz, B.J. (1995) *NDRI*, a locus of *Arabidopsis thaliana* that is required for disease resistance to both a bacterial and a fungal pathogen. *Proceedings of the National Academy of Sciences of the United States of America* 92, 6597-6601.
- 10 Chuck, G., Robbins, T., Nijjar, C., Ralston, E., Courtney-Gutterson, N. and Dooner, H.K. (1993) Tagging and cloning of a petunia flower color gene with the maize transposable element *Activator*. *Plant Cell* 5, 371-378.
- Collins, A., Milbourne, D., Ramsay, L., Meyer, R., Chatot-Balandras, C., Oberhagemann, P., de Jong, W., Gebhardt, C., Bonnel, E. and Waugh, R. (1999)
- 15 QTL for field resistance to late blight in potato are strongly correlated with maturity and vigour. *Molecular breeding* 5, 387-398.
- El-Kharbotly, A., Jacobs, J.M.E., te Lintel-Hekkert, B., Jacobsen, E., Ramanna, M.S., Stiekema, W.J. and Pereira, A. (1996a) Localization of *Ds*-transposon containing T-DNA inserts in the diploid transgenic potato: linkage to the *R1* resistance gene against
- 20 *Phytophthora infestans* (Mont.) de Bary. *Genome* 39, 249-257.
- El-Kharbotly, A., Jacobsen, E., Stiekema, W.J. and Pereira, A. (1995) Genetic localisation of transformation competence in diploid potato. *Theoretical and Applied Genetics* 91, 557-562.
- El-Kharbotly, A., Leonards-Schippers, C., Huigen, D.J., Jacobsen, E., Pereira, A.,
- 25 Stiekema, W.J., Salamini, F. and Gebhardt, C. (1994) Segregation analysis and RFLP mapping of the *R1* and *R3* alleles conferring race-specific resistance to *Phytophthora infestans* in progeny of dihaploid potato parents. *Molecular and General Genetics* 242, 749-754.
- El-Kharbotly, A., Pereira, A., Stiekema, W.J. and Jacobsen, E. (1996b) Race specific
- 30 resistance against *Phytophthora infestans* in potato is controlled by more genetic factors than only *R*-genes. *Euphytica* 90, 331-336.
- Ellis, J. and Jones, D. (1998) Structure and function of proteins controlling strain-specific pathogen resistance in plants. *Current opinion in plant biology* 1, 288-293.

- English, J., J. and Jones, J.D.G. (1998) Epigenetic instability and trans-silencing interactions associated with an SPT::Ac T-DNA locus in tobacco. *Genetics* . Jan. **148**, 457-469.
- 5 Gilliland, L.U., McKinney, E.C., Asmussen, M.A. and Meagher, R.B. (1998) Detection of deleterious genotypes in multigenerational studies. I. Disruptions in individual *Arabidopsis* actin genes. *Genetics* **149**, 717-725.
- Glazebrook, J., Rogers, E.E. and Ausubel, F.M. (1996) Isolation of *Arabidopsis* mutants with enhanced disease susceptibility by direct screening. *Genetics* **143**, 973-982.
- 10 Glazebrook, J., Zook, M., Mert, F., Kagan, I., Rogers, E.E., Crute, I.R., Holub, E.B., Hammerschmidt, R. and Ausubel, F.M. (1997) Phytoalexin-deficient mutants of *Arabidopsis* reveal that *PAD4* encodes a regulatory factor and that four *PAD* genes contribute to downy mildew resistance. *Genetics* **146**, 381-392.
- Graham, K.M. (1963) Inheritance of resistance to *Phytophthora infestans* in two diploid Mexican *Solanum* species. *Euphytica* **12**, 35-40.
- 15 Hammond-Kosack, K.E. and Jones, J.D.G. (1994) Incomplete dominance of tomato *Cf* genes for resistance to *Cladosporium fulvum*. *Molecular Plant-Microbe Interactions* **7**, 58-70.
- Hammond-Kosack, K.E. and Jones, J.D.G. (1997) Plant disease resistance genes. *Annual Review of Plant Physiology and Plant Molecular Biology* **48**, 575-607.
- 20 Hanks, S.K., Quinn, A.M. and Hunter, T. (1988) The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science, USA* **241**, 42-52.
- Innes, R.W. (1998) Genetic dissection of R gene signal transduction pathways. *Current Opinion in Plant Biology* **1**, 299-304.
- 25 Jia, Y., Loh, Y.T., Zhou, J. and Martin, G.B. (1997) Alleles of *Pto* and *Fen* occur in bacterial speck-susceptible and fenthion-insensitive tomato cultivars and encode active protein kinases. *Plant Cell* **9**, 61-73.
- Jones, D.A., Thomas, C.M., Hammond-Kosack, K.E., Balint-Kurti, P.J. and Jones, J.D.G. (1994) Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging. *Science* **266**, 789-793.
- 30 Jones, J.D.G., Romeis, T., Thomas, C., Piedras, P., Durrant, W., Noel, L., Parker, J., Rivas, S., Golstein, C., Dixon, M. and van der Biezen, E. (1999) Molecular genetics of *Cf*- and *RPP*- disease resistance gene function. In *9th International Congress on Molecular Plant-Microbe Interactions*, Biology of Plant-Microbe Interactions, Volume 2 (Wit,

- P.J.G.M.d., Bisseling, T. and Stiekema, W.J., eds). Amsterdam: International Society of Molecular Plant-Microbe Interactions, pp. 243-248.
- Jørgensen, J.H. (1996) Effect of three suppressors on the expression of powdery mildew resistance genes in barley. *Genome* 39, 492-498.
- 5 Kreike, C.M., de Koning, J.R.A., Vinke, J.H., van Ooijen, J.W. and Stiekema, W.J. (1994) Quantitatively inherited resistance to *Globodera pallida* is dominant by one major locus in *Solanum spegazzinii*. *Theoretical and Applied Genetics* 88, 764-769.
- Lawrence, G.J., Finnegan, E.J., Ayliffe, M.A. and Ellis, J.G. (1995) The *L6* gene for flax rust resistance is related to the Arabidopsis bacterial resistance gene *RPS2* and the  
10 tobacco viral resistance gene *N*. *Plant Cell* 7, 1195-1206.
- Leister, R.T., Ausubel, F.M. and Katagiri, F. (1996) Molecular recognition of pathogen attack occurs inside of plant cells in plant disease resistance specified by the Arabidopsis genes *RPS2* and *RPM1*. *Proceedings of the National Academy of Sciences of the United States of America* 93, 15497-15502.
- 15 Leonards-Schippers, C., Gieffers, W., Salamini, F. and Gebhardt, C. (1992) The *R1* gene conferring race-specific resistance to *Phytophthora infestans* in potato is located on potato chromosome V. *Molecular and General Genetics* 233, 278-283.
- Leonards-Schippers, C., Gieffers, W., Schäfer-Pregl, R., Ritter, E., Knapp, S.J., Salamini, F. and Gebhardt, C. (1994) Quantitative resistance to *Phytophthora infestans*  
20 in potato: a case study for QTL mapping in an allogamous plant species. *Genetics* 137, 67-77.
- Liu, Y.G. and Whittier, R.F. (1995) Thermal Asymmetric Interlaced PCR: Automatable Amplification and Sequencing of Insert End Fragments from P1 and YAC Clones to Chromosome Walking. *Genomics* 25, 674-681.
- 25 Martin, G.B., Brommonschenkel, S.H., Chunwongse, J., Frary, A., Ganai, M.W., Spivey, R., Wu, T.Y., Earle, E.D. and Tanksley, S.D. (1993) Map-based cloning of a protein kinase gene conferring disease resistance in tomato. *Science Washington* 262, 1432-1436.
- Mattheij, W.M., Eijlander, R., de Koning, J.R.A. and Louwes, K.M. (1992) Interspecific  
30 hybridization between the cultivated potato *Solanum tuberosum* ssp. *tuberosum* L. and the wild species *Solanum circaeifolium* ssp. *circaeifolium* Bitter exhibiting resistance to *Phytophthora infestans* (Mont.) de Bary and *Globodera pallida* (Stone) Behrens: 1. Somatic hybrids. *Theoretical and Applied Genetics* 83, 459-466.



- McDowell, J.M., Cuzick, A., Can, C., Beynon, J., Dangl, J.L. and Holub, E.B. (2000) Downy mildew (*Peronospora parasitica*) resistance genes in *Arabidopsis* vary in functional requirements for NDR1, EDS1, NPR1 and salicylic acid accumulation. *Plant journal* **22**, 523-529.
- 5    Parker, J.E., Holub, E.B., Frost, L.N., Falk, A., Gunn, N.D. and Daniels, M.J. (1996) Characterization of *eds1*, a mutation in *Arabidopsis* suppressing resistance to *Peronospora parasitica* specified by several different *RPP* genes. *Plant Cell* **8**, 2033-2046.
- 10    Parniske, M., Hammond-Kosack, K.E., Golstein, C., Thomas, C.M., Jones, D.A., Harrison, K., Wulff, B.B.H. and Jones, J.D.G. (1997) Novel disease resistance specificities result from sequence exchange between tandemly repeated genes at the Cf-4/9 locus of tomato. *Cell Cambridge* **91**, 821-832.
- Pereira, A., Aarts, M., van Agtmaal, S., Stiekema, W.J. and Jacobsen, E. (1991) Waxy variegation in transgenic potato. *Maydica* **36**, 323-327.
- 15    Pereira, A. and Aarts, M.G.M. (1998) Transposon tagging with the *En-I* system. In *Arabidopsis protocols*, Methods in Molecular Biology, Volume 82 (Martinez-Zapater, J.M. and Salinas, J., eds). Totowa, New York: Humana Press Inc., pp. 329-338.
- Pereira, A., Jacobs, J.M.E., te Lintel-Hekkert, W., Rutgers, E., Jacobsen, E. and Stiekema, W.J. (1992) Towards the isolation of resistance genes by transposon targeting
- 20    in potato. *Netherlands Journal of Plant Pathology* **98**, 215-221.
- Ritter, E., Debener, T., Barone, A., Salamini, F. and Gebhardt, C. (1991) RFLP mapping on potato chromosomes of two genes controlling extreme resistance to potato virus X (PVX). *Molecular and General Genetics* **227**, 81-85.
- Rommens, C.M.T., Haaren, M.J.J.v., Buchel, A.S., Mol, J.N.M., Tunen, A.J.v.,
- 25    Nijkamp, H.J.J., Hille, J., Van Haaren, M.J.J. and Van Tunen, A.J. (1992) Transactivation of Ds by Ac-transposase gene fusions in tobacco. *Molecular and General Genetics* **231**, 433-441.
- Roupe van der Voort, J., Lindeman, W., Folkertsma, R., Hutten, R., Overmars, H., van der Vossen, E., Jacobsen, E. and Bakker, J. (1998) A QTL for broad-spectrum
- 30    resistance to cyst nematode species (*Globodera* spp.) maps to a resistance gene cluster in potato. *Theoretical and Applied genetics* **96**, 654-661.
- Rudenko, G.N., Nijkamp, H.J.J. and Hille, J. (1994) DS read-out transcription in transgenic tomato plants. *Molecular and General Genetics* **243**, 426-433.

- Salmeron, J.M., Oldroyd, G.E.D., Rommens, C.M.T., Scofield, S.R., Kim, H.-S., Lavelle, D.T., Dahlbeck, D. and Staskawicz, B.J. (1996) Tomato *Prf* is a member of the leucine-rich repeat class of plant disease resistance genes and lies embedded within the *Pto* kinase gene cluster. *Cell* 86, 123-133.
- 5 Song, W.-Y., Wang, G.-L., Chen, L.-L., Kim, H.-S., Pi, L.-Y., Holsten, T., Gardner, J., Wang, B., Zhai, W.-X., Zhu, L.-H., Fauquet, C. and Ronald, P. (1995) A receptor kinase-like protein encoded by the rice disease resistance gene, *Xa21*. *Science* 270, 1804-1806.
- 10 Stam, M., Viterbo, A., Mol Joseph, N.M. and Kooter Jan, M. (1998) Position-dependent methylation and transcriptional silencing of transgenes in inverted T-DNA repeats: Implications for posttranscriptional silencing of homologous host genes in plants. *Molecular and Cellular Biology*. Nov. 18, 6165-6177.
- Tarchini, R., Biddle, P., Wineland, R., Tingey, S. and Rafalski, A. (2000) The complete sequence of 340 kb of DNA around the rice *Adh1-Adh2* region reveals interrupted
- 15 colinearity with maize chromosome 4. *Plant Cell* 12, 381-391.
- Toxopeus, H.J. (1958) Note on the variation in expression of hypersensitivity to *Phytophthora infestans* in potato seedlings. *Euphytica* 7, 38-40.
- Triglia, T., Peterson, M.G. and Kemp, D.J. (1988) A procedure for *in vitro* amplification of DNA segments that lie outside boundaries of known sequences. *Nucleic Acids*
- 20 *Research* 16, 8186.
- Tsugeki, R., Kochieva, E.Z. and Fedoroff, N.V. (1996) A transposon insertion in the Arabidopsis SSR16 gene causes an embryo-defective lethal mutation. *Plant Journal* 10, 479-489.
- Uijtewaal, B.A., Suurs, L.C.J.M. and Jacobsen, E. (1987) Protoplast fusion of
- 25 monohaploid ( $2n = x = 12$ ) potato clones: identification of somatic hybrids using malate dehydrogenase as a biochemical marker. *Plant Science* 51, 277-284.
- Van Engelen, F.A., Molthoff, J.W., Conner, A.J., Nap, J.-P., Pereira A. and Stiekema, W.J. (1995) pBINPLUS: an improved plant transformation vector based on pBIN19. *Transgenic Res.* 4, 288-290.
- 30 Warren, R.F., Merritt, P.M., Holub, E. and Innes, R.W. (1999) Identification of three putative signal transduction genes involved in *R* gene-specified disease resistance in Arabidopsis. *Genetics* 152, 401-412.

- Wastie, R.L. (1991) Breeding for resistance. In *Phytophthora infestans, the cause of late blight in potato*, Advances in Plant Pathology, Volume 7. London: Academic Press, pp. 193-224.
- Whitham, S., Dinesh-Kumar, S.P., Choi, D., Hehl, R., Corr, C. and Baker, B. (1994)  
5 The product of the tobacco mosaic virus resistance gene *N*: similarity to Toll and the Interleukin-1 receptor. *Cell* 78, 1101-1115.
- Wolters, A.M.A., Schoenmakers, H.C.H., van der Meulen-Muisers, J.J.M., van der Knaap, E., Derks, F.H.M., Koornneef, M. and Zelcer, A. (1991) Limited DNA  
elimination from the irradiated potato parent in fusion products of albino *Lycopersicon*  
10 *esculentum* and *Solanum tuberosum*. *Theoretical and Applied Genetics* 83, 225-232.
- Zhou, J., Loh, Y.-T., Bressan, R.A. and Martin, G.B. (1995) The tomato gene *Pti1* encodes a serine/threonine kinase that is phosphorylated by Pto and is involved in the hypersensitive response. *Cell Cambridge* 83, 925-935.
- Zhou, J.M., Tang, X.Y. and Martin, G.B. (1997) The Pto kinase conferring resistance to  
15 tomato bacterial speck disease interacts with proteins that bind a cis-element of pathogenesis-related genes. *EMBO Journal* 16, 3207-3218.


## CLAIMS—

1. An isolated DNA sequence which encodes a protein having the amino acid sequence given in SEQ ID NO: 2 or a functionally homologous protein having an amino acid  
5 sequence showing an identity of at least 80% to SEQ ID NO: 2, which protein confers *Phytophthora infestans* resistance on plants.
2. The DNA sequence according to claim 1, characterized in that the functionally homologous protein has an amino acid sequence showing an identity of at least 85%  
10 to SEQ ID NO: 2.
3. The DNA sequence according to claim 1 or 2, characterized in that it comprises the nucleotide sequence selected from the group consisting of:  
a) the DNA sequence given in SEQ ID NO: 1 and its complementary strand, and  
15 b) DNA sequences hybridizing to the sequences in (a) under stringent hybridization conditions.
4. The DNA sequence according to claim 3, characterized in that it comprises the nucleotide sequence of nucleotides 20 to 1053 of SEQ ID NO: 1.  
20
5. The DNA sequence according to claim 3, characterized in that it comprises the nucleotide sequence of nucleotides 20 to 583 and 727 to 1053 of SEQ ID NO: 1.
6. A protein encoded by the DNA sequence of any of claims 1 to 5.  
25
7. A recombinant vector comprising a DNA sequence under control of an appropriate promoter and regulatory elements for expression in a host cell, wherein the DNA sequence is as defined in any of claims 1 to 5.
- 30 8. Use of a DNA sequence of any of claims 1 to 5 or a recombinant vector of claim 7 for the production of a transgenic plant.
9. A host cell comprising a DNA sequence of any of claims 1 to 5 or a recombinant vector of claim 7.

10. A host cell according to claim 9 which is a plant cell.
11. A plant or any part thereof comprising a plant cell according to claim 10.
- 5 12. Seed, selfed or hybrid progeny or descendant of a plant according to claim 11, or any part thereof.
13. A method of conferring *Phytophthora infestans* resistance on a plant, comprising the steps of
  - 10 i) introducing a DNA sequence of any of claims 1 to 5 or a recombinant vector of claim 7 into a cell of the plant or an ancestor thereof,
  - ii) regenerating plants from the obtained transgenic cells, and
  - iii) selecting plants exhibiting *Phytophthora infestans* resistance.

Figure. 1

1/11



2/11

Fig. 2



3/11





Fig. 3




4/11

Fig. 4



5/11

A Figure 5



B

Pto	-----MGSKYSKATNSINDALSSSYLVPFESYRVPLVD	33
Pti1	MSCFSCDDDDMHRAFDNGPFFMAHNSAGNNGGQRATESAQRETQTVNIQPIAVPSIAVDE	60
StPK-B	-----IKTHQLVSYHE	11
StPK-A	-----IKTNQLISYHE	11
Xa21 kinase	-----	
* * I * * II * * III		
Pto	LEEATNNFDH-KFLIGHGVFGKVKVGLRDGAKVALKRRTPESSQGIIEEFETEITLSFC	92
Pti1	LKDITDNFGS-KALIGEGSYGRVYHGVLSGSGRAAAIKFLD-SSKQPDREFLAQVSMVSR	118
StPK-B	IQQATNNFDDKSNLIGEGSSGSVYKGLSIGTVVAIKVLDLENEQVCKRFDECKVMNRV	71
StPK-A	IQQVTNNFDG-SNLIGEGSSGSVYKGLSSGTTVAIKVLDLENEQVCKRFXTCEVMNRV	70
Xa21 kinase	-----FAPTNNLLSGSGSFGSVYKGLNIQDHSVAVKVLKLENPKALKSFTAECALRM	52
IV V		
Pto	RHPHLVSLIGFCDERNEM-----ILIYKYMENGNLKRHLYG-----SDLPTMSMSWEQRL	142
Pti1	KDENVVELLGYCVDGGFR-----VLAYEYAPNGSLHDILGRKGKVGQAQPGPVLSWAQRV	173
StPK-B	RHRNLVPVITTCSSDYIR-----GFVMPIMPNGSLENWLYKED-----RHLNLHQRV	118
StPK-A	RHRNLVPVITTCSSDYXX-----AFVLKYSXGSHENWLYREV-----RHLNLQRV	117
Xa21 kinase	RHRNLVKIVTICSSIDNRGNDFKAIVYDFMPNGSLEDWIHPETN--DQADQRHLNLHRRV	110
VI * * VII		
Pto	EICIGAARGHLHYLHTRA---IIHRDVKSINILLDENFVPKITDFGISKKGTELDQTHLST	199
Pti1	KIAVGAAGKLEYLHEKAQPHIIHRDIKSSNILLFDDDVAKIADFOLSNQAPDMAARLHST	233
StPK-B	TVMLDAAMAVEYLHHCNVAPIVHCDLKPANVLLDEDMVAHVGDGFGISKILAIKSMAYTE	178
StPK-A	TVMLDAAMAIEYLHHCNDTVIVHCDINPANVLLDEDMVAHVGDGFGISKILAASKSLTQTE	177
Xa21 kinase	TILLDVACALDYLHRRGPEPVVHCDIKSSNVLLDSMDVAHVGDGFGLARILVDGTSLIQOS	170
VIII * IX*		
Pto	V----VKGTGLGYIDPEYFIKGRLTEKSDVYSFGVVLFEVLCARSAIVQSLPREMVNLAEW	255
Pti1	R----VLGTFGYHAPYAMTGQLSSKSDVYSFGVVLLELLTGRKPVVDHTLPRGQQSLVTW	289
StPK-B	T----LGTGLGYIAPEYGSSEIVSASGDVYSYGIMLMEVLTKRPTDEDICNENLDLRKW	233
StPK-A	T----LGTGLGYIAPEYGSSEIVSASGDVYSYGIMLMEVLTKR-----	216
Xa21 kinase	TSSMGFIGTIGYAAPEYGVGLIASTHGDIYSYGILVLEIVTGKRPTDSTFR-PDLGLRQY	229
X XI *		
Pto	AVESHNNQGLEQIVDPNLADKIR-----PESLRKFGDTAVKCLALSSSEDRP	301
Pti1	ATPRLSEDKVKQCVDA RLNTDYP-----PKAIKMAAVALCVQYEADFRP	335
StPK-B	ITQSFS-GSMDVVDANLFSEEEQITCKS-----EMCIASMIELALDCTKKMPESRV	284
StPK-A	-----	
Xa21 kinase	VELGLH-GRVTDVVDTKLILDSENWLNSTNNSPCRRITECIVWLLRLGLSCSQELPSSRT	288
Pto	SMGDVLWKLEYALRLQESVI	321
Pti1	NMSIVVKALQPLLPRPVPS-	354
StPK-B	TMKEVVKRLNLIK-----	297
StPK-A	-----	
Xa21 kinase	PTGDIIDEL-----	297

FIG 6(A)  
SEQ ID NO:1

6/11

TCAGGTGTTG	GTGCTGCAGA	TCAAAACTCA	TCAATTAGTT	TCTTATCAG	50
AGATTCAACA	AGCAACAAAT	AATTTTGATG	ATAAATCAAA	TTTAATTGGT	100
GAGGGAAGCT	CTGGCTCTGT	GTACAAAGGC	ATTTTATCTA	TTGGAAGTGT	150
AGTGGCCATA	AAGGTTCTGG	ATTTGGAAAA	TGAGCAAGTA	TGCAAGAGGT	200
TTGATACCGA	ATGCAAAGTG	ATGAGAAATG	TTAGACACAG	AAATCTTGTT	250
CCAGTGATCA	CTACATGTTT	TAGTGACTAT	ATAAGAGGCT	TTGTTATGCC	300
AATTATGCCC	AATGGAAGTC	TTGAGAATTG	GCTGTACAAA	GAAGATCGCC	350
ACTTGAACCT	TCATCAAAGA	GTAAGTGTA	TGCTTGATGC	AGCTATGGCA	400
GTTGAATATC	TACATCATTG	TCATGTTGCT	CCAATAGTTC	ATTGCGACCT	450
AAAGCCAGCC	AACGTTCTTT	TGGATGAAGA	TATGGTGGCT	CATGTTGGTG	500
ATTTTGGAAT	CTCTAAAATT	TTAGCTATAA	GCAAGTCCAT	GGCCTATACC	550
GAGACATTGG	GTAATCTTGG	ATACATTGCA	CCAGGTATAA	AAAATCTACC	600
CTCTTTGATT	TTCTCTTATC	ATAATTAAAC	CTCTCTAAAT	TCTACCAGTA	650
AGAAAAAGCA	AGGATTTATT	TATGCAGAAT	TATTGTTGTA	TTTCAATTGA	700
GTAACTTTTT	TTCAATTCCT	TTCTAAGAAT	ATGGCTCGGA	GGGAATAGTG	750
TCCGCTAGTG	GTGATGTTTA	TAGTTATGGC	ATTATGTTGA	TGGAGGTTTT	800
GACCAAAAGA	CGGCCAACAG	ATGAAGATAT	ATGCAATGAA	AATCTTGACC	850
TGAGGAAATG	GATAACACAA	TCATTTTCAG	GGAGTATGAT	GGATGTTGTG	900
GATGCTAATC	TTTTTTCTGA	GGAAGAACAA	ATTACTTGTA	AAAGTGAAAT	950
GTGCATAGCC	TCCATGATAG	AATTGGCTTT	AGACTGCACA	AAGAAAATGC	1000
CAGAATCAAG	AGTAACCATG	AAAGAAGTAG	TCAAGAGGCT	TAACAAAATC	1050
AAGAACACAT	TTTTGGAAAT	GTAGAAGTGA	TCAGCATCTC	TTTCTGATCT	1100
GCAAGTTAAC	TTGTTGCTTT	TTGTTTACTG	GTTTCTTTAG	TAAAGGCGTA	1150
TGTACTACTC	GAAGTCATGT	ATTGTTTATA	CTTTAGAGTG	TTGCATTTTG	1200
GAGAAGAAAG	GCATTGTTCC	GAGGAAGTGG	TAATATATCA	TCTCTTTATA	1250
GGTTGGTTGG	TGCAATTGAT	TTTTTAGATT	ATTTTCTATA	AATTTGCGCTC	1300
ACTTGTTTCG	1309				

7/11

Fig. 6(B)

SEQ ID NO:2

IKTHQLVSYHEIQQATNNFDDKSNLIGEGSSGSVYKGILSIGTVVAIKVL  
DLENEQVCKRFDTECKVMRNVRHRNLVPVITTCSSDYIRGFVMPIMPNGS  
LENWLYKEDRHLNLHQVRTVMLDAAMAVEYLLHHCHVAPIVHCDLKPANVL  
LDEDMVAHVGDFGISKILAIKSMAYTETLGTLYIAPEYGSEGIVSASG  
DVYSYGIMLMEVLTKRRPTDEDICNENLDLRKWITQSFSGSMMDVVDANL  
FSEEEQITCKSEMCIASMIELALDCTKKMPESRVTMKEVVKRLNLIK

Fig. 7

8/11

Seq ID NO:3

StPK - A

GGTGTGGTG	CTGGAGATCA	AACTAATCA	ATTGATTTCT	TATCATGAGA	50
TTCAACAAGT	AACAAATAAT	TTTGATGGAT	CCAATTTAAT	TGGCGAGGGA	100
AGCTCTGGCT	CTGTGTACAA	AGGCACATTA	TCAAGTGGAA	CTACGGTGGC	150
CATAAAGGTT	CTGGATTTGG	AAAATGAGCA	AGTATGCAAG	AGGTTTGRTA	200
CAGAATGCGA	AGTGATGAGA	AATGTCAGAC	ATAGAAATCT	TGTTCCAGTG	250
ATTACTACTT	GTTCTAGTGA	CTATAYARCA	GCCTTTGTYC	TGAAATATAT	300
GTCAWATGGG	AGTCACGAAA	ATTGGTTGTA	CAGAGAAGTT	CGCCACTTGA	350
ACCTTCTTCA	AAGAGTCACT	GTAATGCTTG	ATGCGGCTAT	GGCAATTGAA	400
TATCTACATC	ATGGCAATGA	CACTGTGATA	GTTCAATTGCA	GACATAAACC	450
CAGCCAACGT	TCTTTTGGAT	GAAGATATGG	TGGCGCATGT	AGGAGATTTT	500
GGAACTCTTA	AGATCTTAGC	CGCAAGCAAG	TCCCTGACAC	AAACCGAGAC	550
ATTGGGCACT	CTTGGATACA	TTGCACCAGG	TATACTAAAA	TTATAACCTT	600
TCTATTTAAT	TTTTCTCTTA	TCAAAATCAA	GCCCTTGAAA	ATTCTAGGAC	650
TAAATAAAAA	GCAAGTCTTT	GTTAGTATGA	GCATTATTGC	TATATCCAAA	700
TGAGTTAGTT	CTTTTTCATT	TTCTTCTTT	TAAGAGTACG	GCTCAGAAGG	750
AATAGTGTCG	GCTAGTGGTG	ATGTTTACAG	TTACGGCATC	ATGTTGATGG	800
AGGTTTTGAC	GAAAAGAAGG	820			

Seq ID NO:4

StPK - C

ACATTGGGCA	CTCTTGGATA	CATTGCACCA	GGTATAAAAA	ATCTACTCTT	50
TTTATCATAA	TAAAGCCTCT	CCAAATTCTA	CAAGTACAAA	AGCAAGCTTT	100
TATTTATGCA	GAATTATTGT	TGTATTTCAA	TTGAATTAAC	TTTTTTTTCA	150
ATCCTTTTTT	AAGAATATGG	CTCGGATGGA	ATAGTGTCTG	CTAGTGGCGA	200
TGTTTATAGT	TACGGCATCA	TGTTGATGGA	GGTTTTGACG	AAAAGAAGGC	250
CAACAAATGA	AGAGATATGC	AATGAAAATC	TTGACTTGAG	GAAATGGATC	300
ACACAATCAT	TTTCAGGGAG	TATGATGGAC	GTTGTGGATG	CCAATCTTTT	350
CTCCGAGGAA	GAACAGATCA	CTTCAGAAAG	TGAAATCTGC	ATAGCGTCCA	400
TGATAGAATT	GGGTTTAGAC	TGCACAAAGA	AAATGCCAGA	ATCAAGA	447

Seq ID NO:5

StPK - D

AACATTGGGC	ACTCTTGGAT	ACATTGCACC	AGGTATACTT	AAATTATAAC	50
CTATCTATTT	GATTTTCTC	TTATCAAAAT	CAAGCCCTTG	AAAATTCTAG	100
GACTAAATAA	AAAGCAAGTC	TTTGTATTAT	GTACAAGCAT	TATTGTTATA	150
TCCAAATGAG	TTATTCTTTT	TCATTTTCGA	TCTTTTAAGA	ATATGGCTCA	200
GAAGGAATAG	TTTCCGCTAG	TGGTGATGTT	TACAAGGACT	GTGATGGACG	250
TTGTGGATTG	CAACCTTTTT	TGTGAGGAAG	AACAAATCAC	TAGGAAAAGT	300
GAAATCTGCA	TAGCTCCAT	GATAGAATTG	GCCTTAGATT	GCACAAAGAA	350
AATGCCAGAA	TCAAGA	366			

Seq ID NO:6

StPK - E

ACATTGGGCA	CTCTTGGATA	CATGGCACCA	GGTATAAAAA	AGAATCTACT	50
------------	------------	------------	------------	------------	----

Fig.7 (cont.)

9/11

CTCTTTGATT	TTCTCTTATC	ATAATTAATT	AAGCCTCTCC	AAGTTCTAGA	100
AGTAAAAGAT	GCAAGTTTTT	ATTTATTGAG	AATTATTGTT	GTATTTCAAT	150
TGAATAACTG	TTTTTTTCTC	AACCCTTTTC	TATGAATATG	GCTCGGAGGG	200
AATAGTGTCC	ACTAGTGGTG	ATGTTTATAG	TTACGGCATC	ATGTTGATGG	250
AGGTTTTGAC	CAAGAGAAGG	CCAACAGATG	AAGAGATATG	CAATGAAACT	300
CTTGACTTGA	GGAAATGGAT	CACACAATCA	TTTTTCAGGA	GTATGATGGA	350
CGTTGTGGAT	GCCAATCTTT	TCTCCGAGGA	AGAACAGATC	ACTTCAGAAA	400
GTGAAATCTG	CATTGCGTCC	ATGATAGAAT	TGGGTCTAGA	CTGCACAAAG	450
AAAATGCCAG	AATCAAGA	468			

Seq ID NO:7

StPK - F

AAAAAAGCAA	GTCTTCATTT	AGGCAGAATT	ATTGTTGTAT	TTCAAGGGAG	50
TAACTTTTCC	TCAATCCTTT	TCTAAGAATA	TGGCTCAGAG	GGAATAGTGT	100
CTTCTAGTGG	TGATGTTTAT	AGCTATGGCA	TCATGTTGAT	GGAAGTCTTG	150
ACCGAAAGAA	GGCCAACAGA	TGAAGAGATA	TGCAATGAAA	ATCTTGACCT	200
GAGGAAATGG	ATAATACAAT	CATTTTCAGG	GAGTATGATG	GACGTTGTCTG	250
ATGCCAATCT	TTTTTACGAG	GAAGAACAAA	CCACTAGTAA	AAGTGAAATC	300
TGCATAGCGT	CCATGATAGA	ATTGGGTTTA	GATTGCACAA	AGAAAATGCC	350
AGAATCAAGA	360				

Seq ID NO:8

StPK - G

TCTCCAAGTT	GTAGAAGTAA	AAAGAGAACT	ATTGTTATAT	TTCAATGAG	50
CAACTTTTGG	TCAATCATTT	TCTAAGAATA	TGGATCAGAG	GGAATAGTGT	100
CTGCTAGTGG	TGATGTTTAT	AGCTACGGAA	TCATGTTGAT	GGAGGTTTGG	150
ACCAAAGAA	GGCCAACAGA	TGAAGAGATA	TGTAATCAAA	ATCTTGACCT	200
GAGGAAATGG	ATAATACAAT	CATTTTCAGG	GAGTATGACG	GACATCGTGG	250
ATGCCAATAT	TTTTTCTGAG	GAAGAACAAA	TTACTTGTA	AAGTAAATG	300
TGCATAGCCT	CCATGATAGA	ATTGGCTTTA	GACTGCACAA	AGAAAATGCC	350
AGAATCAAGA	360				

Seq ID NO:9

StPK - H

ACATTGGGCA	CTCTTGATA	CATTGCACCA	GGTATAAGAA	AATCTACTCT	50
CATTGATTTT	CTCTTATCAT	AATTAAGCCT	CTCCAATTGT	TGTAGAAGTA	100
AAAATAGAAT	CATTGTATTT	CAATTGAGTA	ACCTTTCTTC	AATCCTTTTC	150
TAAGAATATG	GCTCGGAGGG	AATAGTGAAT	GTCTACTAGT	GGTGATGTTT	200
ATAGCTACGG	CATCATGCTG	ATGGAGGTTT	TGACGAAAAG	AAGGCCAACA	250
GATGAAGAGA	TATGCAATGA	AATTCTTGAC	TTGAGGAAAT	GGATCACACT	300
ATCATTTTCA	GGGAGTATGT	TGGACATTGT	GGATGCCAAT	ATTTTTTGTTG	350
AGGAAGAACA	AATCACTAGT	AAAAGTGAAA	TGTGCATAGC	CTCCATGATA	400
GAACCGTCTT	TAGACTGCAC	AAAGAAAATG	CCAGAATCAA	GA	442

Seq ID NO:10

StPK - I

ATTGGGCACT	CTTGGATACA	TAGCACCAGG	TATAAAAAAA	TTTACTCTCT	50
TTGATTTTCT	TTTATATCAT	AATTAAGCCT	CTCCAAATTC	TACAAGTAGA	100
AAAAACAAG	TTTTCATTTA	TGCAGAATTA	TTGTTGTATT	TCAATTGAGT	150
AACTTTTCTT	CAATCCTTTT	CTAAGAATAT	GGCTCAAAGG	GAATAGTGTG	200
TGCTAGTGGT	GATGTTTATA	GCTATGGCAT	CATGTTGACC	TGAGGAAATG	250

10/11

Fig.7 (cont.)

GATAATACAA	TCTTGATGAA	AAGCTATGCA	ATGAAAATCT	TGACCTGAGG	300
AAATGGATAA	TACAATCATT	TTTAGGGAGT	ATGATGGACA	TTGTGGATGC	350
CAATCTTTTT	TGTGAGGAAG	TACAAATCAC	TTGTAAAAGT	GAAATGTGCC	400
TAGCCTCCAT	GATAGAATTG	GCTTTAGATT	GCACAAAGAA	AATGCCAGAA	450
TCAAGA	456				

Seq ID NO:11

StPK - J

ACATTGGGCA	CTCTTGGATA	CATTGCACCA	AGGTATAAAA	AATCTACTCA	50
CTTTGATTTT	CTTTTATCAT	AATAAAGCCT	CTCAAATTC	TACAAGTATA	100
AAAGCAACCT	TTTATTTATG	CAGAATTATT	GTTGTATTTT	AATTGAATTA	150
ACTTTTTTTT	CAATCCTTTT	TTAAGAATAT	GGCTCGGATG	GAATAGTATC	200
TGCTAGTTGC	GATGTTTATA	GTTACGGCAT	CATGTTGATG	GAGGTTTTGA	250
CGAAAAGAAG	GCCAACAGAT	GAAGAGATAT	GCAATGAAAA	TCTTGACCTG	300
AGGAAATGGA	TAATACAATC	ATTTTCAGGG	AGTATGATGG	ACGTTGTCGA	350
TGCCAATCTT	TTTACGAGGA	AGAACAAATC	ACTAGTAAAA	GTGAAATCTG	400
CATAGCGTCC	ATGATAGAAT	TGGGTTTAGA	TTGCACAAAG	AAAATGCCAG	450
AATCAAGA	458				


Seq ID NO:12

StPK - K

ACATTGGGCA	CTCTTGGATA	CATTGCACCA	GGTATAAAAA	AATCTACTCT	50
CTTTGATTTT	CTCTTATATC	ATAATTAAGC	CTCTCTAAGG	TCTAAAAGTT	100
AAAAAAAAAA	AAAAACAAGT	TTTCATTTAT	GCAGAATTAT	TGTTGAATTT	150
CAATTGAGTA	ACTTTTCCTC	AATCCTTCTC	TAAGAATATG	GCTCGGAGGG	200
AATAGTGTCT	GCTAGTGGTG	ATGTTTATAG	CTACGGCATC	ATGTTGATGG	250
AGGTTTTGAC	GAAAAGAAGG	CCAACAGATG	AAGAGATATG	CAATGAAAAT	300
CTTGACTTGA	GGAAATGGAT	CACACAATCA	TTTTTCAGGA	GTATGATGGA	350
TGTTGTGGAT	GCCAATCTAT	TTTCTGCGGA	AGAACAAATC	ACTAGTAAAA	400
GTGAAATGTG	CATAGCCTCC	ATGATAGAAT	TGGCTTTAGA	CTGCACAAAG	450
AAAATGCCAG	AATCAAGA	468			

Figure 8:

11/11





(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number  
**WO 02/038727 A3**

(51) International Patent Classification<sup>7</sup>: C12N 9/12,  
15/82, 15/63

(21) International Application Number: PCT/NL01/00804

(22) International Filing Date:  
5 November 2001 (05.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
00203911.3 9 November 2000 (09.11.2000) EP

(71) Applicant (for all designated States except US): PLANT  
RESEARCH INTERNATIONAL B.V. [NL/NL];  
Droevendaalsesteeg 1, NL-6708 PB Wageningen (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): VAN ENCK-  
EVORT, Leonora, Johanna, Gertruda [NL/NL]; Cort  
van der Lindenstraat 35, NL-6702 BD Wageningen  
(NL). JACOBSEN, Evert [NL/NL]; Rizema Bosweg  
77, NL-6708 DK Wageningen (NL). STIEKEMA,  
Willem, Johannes [NL/NL]; Leonard Roggeveenstraat  
21, NL-6708 NB Wageningen (NL). PEREIRA, Andy  
[NL/NL]; Indira Gandhi Singel 14, NL-6717 GG EDE  
(NL).

(74) Agent: DE HOOP, Eric; Octrooibureau Vriesendorp &  
Gaade, P.O. Box 266, NL-2501 AW The Hague (NL).

(81) Designated States (national): AB, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,

LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,  
ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

**Declaration under Rule 4.17:**

as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB,  
GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP,  
KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,  
MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD,  
SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN,  
YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ,  
SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY,  
KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH,  
CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT,  
SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,  
GQ, GW, ML, MR, NE, SN, TD, TG)

**Published:**

with international search report

(88) Date of publication of the international search report:  
28 November 2002

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: DNA SEQUENCES ENCODING PROTEINS CONFERRING PHYTOPHTHORA INFESTANS RESISTANCE ON  
PLANTS

(57) Abstract: Genomic sequences encoding *Phytophthora infestans* resistance proteins are provided herein. Specifically, sequences  
from potato required for *P. infestans* resistance have been cloned and sequence provided, together with the encoded amino acid se-  
quence. DNA encoding the amino acid sequence or amino acid sequences showing a significant degree of homology thereto may be  
introduced into plant cells and the encoded polypeptide expressed, conferring *P. infestans* resistance on plants comprising such cells  
and descendants thereof.

WO 02/038727 A3

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/NL 01/00804

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/12 C12N15/82 C12N15/63

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, SEQUENCE SEARCH, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	VAN ENCKEVORT L.J.G.: "Identification of potato genes involved in Phytophthora infestans resistance by transposon mutagenesis (thesis)" 21 November 2000 (2000-11-21), WAGENINGEN UNIVERSITY, NL XP002210496 ISBN: 90-5808-343-8 page 85, paragraph 3 -page 93, paragraph 4	1-13
A	VAN ENCKEVORT L.J.G. ET AL.: "Selection of independent Ds transposon insertions in somatic tissue of potato by protoplast regeneration" THEORETICAL AND APPLIED GENETICS, vol. 101, September 2000 (2000-09), pages 503-510, XP000926420 the whole document	1-13

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

21 August 2002

Date of mailing of the international search report

10/09/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo.nl,  
Fax (+31-70) 340-3018

Authorized officer

Schönwasser, D

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 01/00804

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EL-KHARBOTLY A. ET AL.: "Localization of Ds-transposon containing T-DNA inserts in the diploid transgenic potato: linkage to the R1 resistance gene against <i>Phytophthora infestans</i> (Mont.) de Bary" GENOME, vol. 39, no. 2, April 1996 (1996-04), pages 249-257, XP000926428 cited in the application the whole document	1-13
A	TARCHINI R. ET AL.: "The Complete Sequence of 340 kb of DNA around the Rice Adh1-Adh2 Region Reveals Interrupted Colinearity with Maize Chromosome 4" THE PLANT CELL, vol. 12, no. 3, March 2000 (2000-03), pages 381-391, XP002168485 cited in the application page 382, column 2, line 10 -page 383, column 2, line 6; table 1 page 389, column 2, paragraph 2	1-6
A	KANEKO T. ET AL.: "Structural analysis of Arabidopsis thaliana chromosome 5.V. Sequence features of the regions of 1,381,565 bp covered by twenty one physically assigned P1 and TAC clones" DNA RESEARCH, vol. 5, no. 1, 30 April 1998 (1998-04-30), pages 131-145, XP000926429 cited in the application page 135: figure legend, second line from the bottom; page 139: line 8 of the first table (MPL12)	1-6
A	DATABASE EMBL 'Online! 28 July 1999 (1999-07-28) ALCALA J. ET AL.: "Generation of ESTs from tomato carpel tissue; EST267901 tomato ovary, TAMU Lycopersicon esculentum cDNA clone cLED34E19, mRNA sequence" Database accession no. AI898458 XP002168486 abstract	1-6

-/--

# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/NL 01/00804

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SUNDARESAN V. ET AL.: "PATTERNS OF GENE ACTION IN PLANT DEVELOPMENT REVEALED BY ENHANCER TRAP AND GENE TRAP TRANSPOSABLE ELEMENTS"</p> <p>GENES AND DEVELOPMENT, vol. 9, no. 14, 15 July 1995 (1995-07-15), pages 1797-1810, XP000674520 ISSN: 0890-9369 the whole document</p>	1-13
A	<p>MEISSNER R. ET AL.: "A new model system for tomato genetics"</p> <p>THE PLANT JOURNAL, vol. 12, no. 6, 1997, pages 1465-1472, XP002096585 ISSN: 0960-7412 the whole document</p>	1-13

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**